



**You have downloaded a document from  
RE-BUS  
repository of the University of Silesia in Katowice**

**Title:** The first detection of the alien species: green-peach aphid *Myzus (Nectarosiphon) persicae* (Insecta, Hemiptera, Aphididae) in the Svalbard archipelago

**Author:** Karina Wieczorek, Dominik Chłond

**Citation style:** Wieczorek Karina, Chłond Dominik.(2019). The first detection of the alien species: green-peach aphid *Myzus (Nectarosiphon) persicae* (Insecta, Hemiptera, Aphididae) in the Svalbard archipelago. "Polar Biology" (Vol. 42, iss. 10 (2019), s. 1947-1951), doi 10.1007/s00300-019-02562-9



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI  
W KATOWICACH



Biblioteka  
Uniwersytetu Śląskiego



Ministerstwo Nauki  
i Szkolnictwa Wyższego



# The first detection of the alien species: green-peach aphid *Myzus (Nectarosiphon) persicae* (INSECTA, HEMIPTERA, APHIDIDAE) in the Svalbard archipelago

Karina Wieczorek<sup>1</sup> · Dominik Chłond<sup>1</sup>

Received: 11 April 2019 / Revised: 13 August 2019 / Accepted: 17 August 2019 / Published online: 28 August 2019  
© The Author(s) 2019

## Abstract

Although the high Arctic archipelago Svalbard is among the best-studied Arctic regions in terms of biodiversity, its aphid fauna is extremely limited. Two endemic species have been described to date, and no species of a foreign origin has been registered. Our observation is the first record of the non-native and potentially invasive green-peach aphid species *Myzus (Nectarosiphon) persicae* (INSECTA, HEMIPTERA, APHIDIDAE), which was collected on the largest island of the archipelago—Spitsbergen. The green-peach aphid was found on three different ornamental plants: *Viola tricolor* (Violaceae), *Dianthus caryophyllus* (Pink Kisses) (Caryophyllaceae) and *Petunia* × *hybrida* (surfinia) (Solanaceae). The plants were displayed in front of one of the shops in the main street of Longyearbyen, the largest populated settlement of Svalbard. The infected plants came from the only supermarket on the archipelago, which had brought them from the mainland of Norway. Although the aphids (winged and wingless viviparous females and immature ones) were clearly visible on all of the plants (the distinct colonies from few to numerous individuals), the owners were not aware that the plants were infected. Similarly, the supermarket staff who are responsible for the live green plants did not register the presence of aphids on the imported plants. In this context, our results also suggest that a stricter approach to regional biosecurity needs to be considered in order to avoid the risks of further unintentional introductions.

**Keywords** High arctic · Biological invasion · Macrosiphini · Pest species

## Introduction

The high Arctic archipelago Svalbard is among the best-studied Arctic regions in terms of biodiversity (Prestrud et al. 2004; Johansen et al. 2012). Despite its isolation, invasions have also occurred in this area and are widely recognised as constituting a serious risk to the Arctic region (Coulson et al. 2013; Wisz et al. 2015). Additional factors are the rapid climate change in the region and increases in human activity (Ware et al. 2011, 2014, 2016). Most of the alien species that have been introduced to Svalbard

are plants. The percentage of alien plant taxa that was recorded in Svalbard (57%) compared to native species is higher than that recorded in other Arctic regions (Alsos et al. 2015). The highest species diversity and abundance of aliens were recorded in or close to the human settlements of Barentsburg, Pyramiden or Longyearbyen on the largest island Spitsbergen (Liška and Soldán 2004; Roalsø 2012). However, in the harsh Arctic conditions, only a few of the introduced plant species have established stable populations (Alsos et al. 2007). The current alien terrestrial invertebrate fauna of Svalbard consists of a relatively few species and none of them have yet displayed invasive tendencies (Coulson 2015).

The aphid fauna of Svalbard is extremely limited and comprises two endemic species—*Acyrtosiphon svalbardicum* (Heikinheimo) (Hodkinson et al. 2002) and *Sitobion calvulus* (Ossiannilsson) (Hodkinson et al. 2004). Non-native species of this group of insects have not as yet been recorded.

✉ Karina Wieczorek  
karina.wieczorek@us.edu.pl

Dominik Chłond  
dominik.chlond@us.edu.pl

<sup>1</sup> Department of Zoology, Faculty of Biology  
and Environmental Protection, University of Silesia,  
Bankowa 9, 40-007 Katowice, Poland

During the investigation, which was conducted in August 2018, we had the opportunity to collect aphids (INSECTA, HEMIPTERA, APHIDIDAE) in Longyearbyen, which is the largest settlement in Svalbard, and its vicinity, as well as in the research town of Ny-Ålesund, both of which are located on the island of Spitsbergen in Svalbard. One of the results of our field study revealed the first detection of the alien aphid species *Myzus (Nectarosiphon) persicae* (Sulzer) in the Svalbard archipelago. The green-peach aphid is a polyphagous generalist that is globally distributed and is listed among the 14 species of the greatest economic importance that has a pest status (van Emden and Harrington 2007).

## Materials and methods

Longyearbyen is the largest populated settlement on the Svalbard archipelago, and is located in the High Arctic, 78° North. Longyearbyen lies on Spitsbergen on the southern side of Adventfjorden, and is stretched out along the Longyear River. The surveys that are reported here were primarily carried out in the town and its vicinity (Fig. 1).

The investigation was conducted for a period of two weeks from 10 to 24 August 2018. The permission for the research application for RiS-ID 10,994 (including aphid collection) was issued by the Governor of Svalbard.

The aphids were collected directly from the host plants with a fine hair brush and placed into Eppendorf tubes containing 70% ethanol. The location, sampling date and host plant name were recorded on the labels that were placed

onto the tubes. Adult wingless or winged females were slide-mounted using the method of Kanturski and Wieczorek (2012) and identified to the species level. The slides were examined using a Nikon Ni-U light microscope equipped with a phase contrast system. Field photographs were taken using an iPhone 7 camera with the OlloClip Macro Pro Lens Set. Samples were identified by K. Wieczorek based on their morphological diagnostic features using the standard literature-based keys (Blackman 2010; Blackman and Eastop 2006, 2018). The aphid material is deposited in the collection of the Department of Zoology, University of Silesia in Katowice, Katowice, Poland (DZUS):

DZUS 23/8.18\_6 *Myzus (Nectarosiphon) persicae*, three alate viviparous females, Longyearbyen, Norway, 23.08.2018, *Viola tricolor*, K.Wieczorek leg.; DZUS 23/8.18\_7 *Myzus (Nectarosiphon) persicae*, two alate viviparous females, one apterous viviparous female, Longyearbyen, Norway, 23.08.2018, *Dianthus caryophyllus* (Pink Kisses), K.Wieczorek leg.; DZUS 23/8.18\_8 *Myzus (Nectarosiphon) persicae*, four alate viviparous females, Longyearbyen, Norway, 23.08.2018, *Petunia* × *hybrida*, K.Wieczorek leg.

## Results and discussion

Living specimens of a virginoparous generation of *Myzus (Nectarosiphon) persicae* were collected from plants displayed in front of one of the shops in the main street of Longyearbyen (78°13'02.77"N 15°38'08.69"E) (Fig. 1).




Fig. 1 Collecting area



Aphids were found on three different decorative plants: *Viola tricolor* (Violaceae) (Fig. 2a), *Dianthus caryophyllus* (Pink Kisses) (Caryophyllaceae) (Fig. 2b) and *Petunia* × *hybrida* (surfinia) (Solanaceae) (Fig. 2c). All of these are annuals or short-lived perennials that are commonly used as ornamental plants. They can tolerate relatively harsh conditions.

The abovementioned plants are secondary hosts for *M. (N.) persicae* as its primary host *Prunus* spp. is absent in Svalbard. The highest density of adult and immature morphs were observed on *Viola tricolor* (on both sides of older leaves, Fig. 2d) and *Dianthus caryophyllus* (Pink Kisses) (under the inflorescence and between the petals, Fig. 2e,

**Fig. 2** The green-peach aphid *Myzus (Nectarosiphon) persicae* and its host plants: **a** *Viola tricolor*; **b** *Dianthus caryophyllus* (Pink Kisses); **c** *Petunia* × *hybrida*; **d, e** winged viviparous female and immature ones; **f** wingless viviparous females and immature ones; **g** winged viviparous female



**Table 1** The number of morphs of the green-peach aphid *Myzus (Nectarosiphon) persicae* observed on its host plants

<i>Myzus (Nectarosiphon) persicae</i>	<i>Viola tricolor</i>	<i>Dianthus caryophyllus</i> (Pink Kisses)	<i>Petunia</i> × <i>hybrida</i>
Number of individuals			
Wingless viviparous females	17	21	
Winged viviparous females	11	13	19
Immature	24	38	

f). *Surfinia* was the least infested—only a small number of winged viviparous females were observed (Fig. 2g). The aphids were clearly visible on all the plants. The exceptions were the immature morphs, which were hidden in the petals of *Dianthus*. The immature morphs and winged morphs had the largest percentage share in the observed colonies (Table 1). Since neither oviparous females nor males (the sexual generation) were found in August and it is lack of its secondary host—*Prunus* spp., this species reproduces by anholocycly (permanent parthenogenesis) in high Arctic conditions of the Svalbard archipelago.

The infected plants came from the only supermarket on the archipelago, which had brought them from the mainland of Norway in June 2018. The plants were purchased also in June, and at the end of August, they were still in good condition with no visible signs of damage despite the aphids that were feeding on them. The owners of the plants were neither aware that the plants were infected nor of the presence of aphids. Similarly, the supermarket staff who are responsible for the live green plants did not register the presence of aphids on the imported plants. Although the average summer temperatures in Longyearbyen are quite low (3–7 °C), both the aphids and plants were in good condition, probably because they were kept in a closed room at night (which is required as polar bears roam through the town during the night). No other species of aphids (native or alien) or additional records of *M. (N.) persicae* in the studied area were found.

Unintentional introductions present the greatest threats to the Arctic and Antarctic ecosystems (Frenot et al. 2005; Snyder and Anions 2008; Chown et al. 2012; Hughes et al. 2015; Chown and Convey 2016; Chan et al. 2018). Alsos et al. (2015) stated that the ornamental plant trade and introduction through agriculture are of limited relevance in the high Arctic—only one observation of potato cultivation and one of the ornamental plant *Viola tricolor* exist, both from Longyearbyen. However, during our brief stay in Longyearbyen, we observed more examples of ornamental plants that were placed in front of shops or hotels and a wide variety of domestic pot plants available at the local supermarket. In this aspect, *M. (N.) persicae*, as a highly polyphagous species that is associated with more than 40 plant families and over 400 species (a secondary hosts) (Blackman and Eastop 2000), has also an advantage in colonising new areas.

Moreover, it can transmit more than 100 plant viruses and its high levels of resistance to almost all classes of insecticides are well recognised (Bass et al. 2014). It has been proved that in similar harsh conditions, the introduced aphids, at least in some cases, prefer native host plants to exotic host plants (Hullè 2012). They can also transfer viruses (e.g. *Barley yellow dwarf virus*) to native plant species (Svanella-Dumas et al. 2013).

Our observation is the first one to record this non-native, potentially invasive and extremely harmful aphid species that has been collected in the archipelago. Whether the recorded population of *M. (N.) persicae* survived in small populations for a short time period and either died out naturally or were removed by human intervention (our recommendation for the owners was to destroy all infested plants) remains a speculation. However, our results suggest that a stricter approach to regional biosecurity needs to be considered in order to avoid the risks of further unintentional introductions. An additional challenge is to increase the public awareness of invasive and alien species of aphids in the fragile ecosystem of Svalbard. Such baseline information is critical for creating the appropriate management strategies.

**Acknowledgements** The research has received funding from the European Union's Horizon 2020 project INTERACT under Grant Agreement No 730938. Our particular thanks to Hannele Savela, Coordinator INTERACT Transnational Access and Christina A. Pedersen, Manager of the Sverdrup Research Station, Norwegian Polar Institute, Ny-Ålesund. We are very grateful to all the staff of Sverdrup Research Station, Norwegian Polar Institute, Ny-Ålesund for help and hospitality during our study. We thank Michelle Simmons for linguistic improvement of the second version of the manuscript. We are also grateful to the Editor and to the anonymous Reviewers for all valuable comments during the review process.

**Author contribution** KW and DCH designed the research, collected the insects and analysed the results. KW wrote the paper. DCH prepared the figures. Both authors have reviewed the paper.

## Compliance with ethical standards

**Conflict of interest** The authors have no conflict of interest to declare.

**Ethical approval** This article does not contain any studies with human participants performed by any of the authors.

**Human and animals rights** All applicable international and national guidelines for the care and use of animals were followed. All proce-



dures performed in studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Alsos IG, Eidesen PB, Ehrich D, Skrede I, Westergaard K, Jacobsen GH, Landvik JY, Taberlet P, Brochmann C (2007) Frequent long-distance plant colonization in the changing Arctic. *Science* 316:1606–1609
- Alsos IG, Ware C, Elven R (2015) Past Arctic alien have passed away, current ones may stay. *Biol Invasions* 17:3113. <https://doi.org/10.1007/s10530-015-0937-9>
- Bass Ch, Puinean AM, Zimmer ChT, Denholm I, Field LM, Foster SP, Gutbrod O, Nauen R, Slater R, Williamson MS (2014) The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*. *Insect Biochem Mol Biol* 51:41–51. <https://doi.org/10.1016/j.ibmb.2014.05.003>
- Blackman RL (2010) Aphids—Aphidinae (Macrosiphini): Handbook for identification of British insects. Royal Entomological Society, St Albans
- Blackman RL, Eastop VF (2000) Aphids on the world's crops: an identification and information guide, 2nd edn. Wiley, Chichester
- Blackman RL, Eastop VF (2006) Aphids on the world's herbaceous plants and shrubs. Wiley, Chichester
- Blackman RL, Eastop VF (2018) Aphids of the world's plants: an online identification and information guide. <https://www.aphidsonworldplants.info>. Accessed 30 Aug 2018
- Chan FT, Stanislawczyk K, Sneekes AC, Dvoretzky A, Gollasch S, Minchin D, David M, Jelmert A, Albretsen J, Bailey SA (2018) Climate change opens new frontiers for marine species in the Arctic: current trends and future invasion risks. *Glob Change Biol* 25:25–38. <https://doi.org/10.1111/gcb.14469>
- Chown SL, Huiskes AHL, Gremmen NJM, Lee JE, Terauds A, Crosbie K, Frenot Y, Hughes KH, Imura S, Kiefer K, Lebouvier M, Raymond B, Tsujimoto M, Ware Ch, Van de Vijver B, Bergstrom DM (2012) Continent-wide risk assessment for the establishment of nonindigenous species in Antarctica. *PNAS* 109:4938–4943. <https://doi.org/10.1073/pnas.1119787109>
- Chown SL, Convey P (2016) Antarctic entomology. *Annu Rev Entomol* 61:119–137. <https://doi.org/10.1146/annurev-ento-010715-023537>
- Coulson SJ et al (2013) Introduction of invertebrates into the high Arctic via imported soils: the case of Barentsburg in Svalbard. *Biol Invasions* 15:1–5
- Coulson SJ (2015) The alien terrestrial invertebrate fauna of the high Arctic archipelago of Svalbard: potential implications for the native flora and fauna. *Polar Res* 34(1):27364. <https://doi.org/10.3402/polar.v34.27364>
- Frenot Y, Chown SL, Whinam J, Selkirk PM, Convey P, Skotnicki M, Bergstrom DM (2005) Biological invasions in the Antarctic—extent, impacts and implications. *Biol Rev* 80:45–72. <https://doi.org/10.1017/S1464793104006542>
- Hodkinson ID, Coulson SJ, Bird J, Webb NR (2002) Discovery of the rare alate morph *Acyrtosiphon svalbardicum* Heikkinheimo (Homoptera: Aphididae): descriptions and implications for species ecology. *Norw J Entomol* 49:77–80
- Hodkinson ID, Bird JM, Cooper EJ, Coulson SJ (2004) The sexual morphs of the endemic Svalbard aphid *Acyrtosiphon calvulus* (Ossiannilsson), with notes on species biology. *Norw J Entomol* 51:131–135
- Hughes KA, Perterra LR, Molina-Montenegro MA, Convey P (2015) Biological invasions in terrestrial Antarctica: what is the current status and can we respond? *Biodivers Conserv* 24(5):1031. <https://doi.org/10.1007/s10531-015-0896-6>
- Hullè M (2012) *Myzus ascalonicus*, an aphid recently introduced to Sub-Antarctic Islands, prefers native to exotic host-plants. *Environ Entomol* 41(6):1398–1404. <https://doi.org/10.1603/EN12057>
- Johansen B, Karlens S, Tømmervik H (2012) Vegetation mapping of Svalbard utilising Landsat TM/ETM data. *Polar Record* 48(1):47–63. <https://doi.org/10.1017/S0032247411000647>
- Kanturski M, Wiczorek K (2012) Mszyce (Homoptera, Aphidoidea) jako grupa modelowa w badaniach zoocenologicznych. In: Kuczer M (ed) Nowe trendy w naukach przyrodniczych. Creative science [in Polish]. CREATIVETIME, Cracow, pp 123–131
- Liška J, Soldán Z (2004) Alien vascular plants recorded from the Barentsburg and Pyramiden settlements, Svalbard. *Preslia* 76:279–290
- Prestrud P, Strøm H, Goldman HV (2004) A catalogue of the terrestrial and marine animals of Svalbard. Norwegian Polar Institute, Trømso
- Roalsø ER (2012) Alien plant species in Svalbard. Norwegian University of Science and Technology, Norwegian, pp 1–25
- Snyder E, Anions M (2008) Pathways analysis of invasive plants and insects in the northwest territories. Report for the Department of Environment and Natural Resources, Wildlife Division, Government of the Northwest Territories. Project No: PM 005529:1–28
- Svanella-Dumas L, Candresse T, Hullè M, Marais M (2013) Distribution of *Barley yellow dwarf virus*-PAV in the Sub-Antarctic Kerguelen Islands and characterization of two new *Luteovirus* species. *PLoS ONE* 8(6):e67231. <https://doi.org/10.1371/journal.pone.0067231>
- van Emden F, Harrington R (2007) Aphids as crop pests. CABI Publishing, London
- Wisz SM, Broennimann O, Grønckjaer P, Moller PR, Olsen SM, Swingedouw D, Hedeholm RB, Nielsen EE, Guisan A, Pellissier L (2015) Arctic warming will promote Atlantic-Pacific fish interchange. *Nat Clim Change* 5:261–265
- Ware Ch, Bergstrom DM, Müller E, Alsos IG (2011) Humans introduce viable seeds to the Arctic on footwear. *Biol Invasions* 14(3):567–577. <https://doi.org/10.1007/s10530-011-0098-4>
- Ware Ch, Berge J, Sundet JH, Kirkpatrick JB, Coutts ADM, Jelmert A, Olsen SM, Floerl O, Wisz MS, Alsos AG (2014) Climate change, non-indigenous species and shipping: assessing the risk of species introduction to a high-Arctic archipelago. *Divers Distrib* 20:10–19. <https://doi.org/10.1111/ddi.12117>
- Ware Ch, Berge J, Jelmert A, Olsen SM, Pellissier L, Wisz M, Kriticos D, Semenov G, Kwaśniewski S, Alsos IG (2016) Biological introduction risks from shipping in a warming Arctic. *J Appl Entomol* 53:340–349

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.